

Fast, Simple QuEChERS Extraction and Cleanup of Pesticide Residue Samples

- Achieve a four-fold increase in sample throughput.
- Significantly reduce material costs.
- Convenient, ready to use centrifuge tubes with ultra pure, pre-weighed adsorbent mixtures.



Quick, Easy, Cheap, Effective, Rugged, and Safe, the QuEChERS (“catchers”) method for extracting pesticides from food is based on research by the US Department of Agriculture.¹ In addition to using less solvent and materials versus conventional SPE methods, QuEChERS employs a novel and much quicker dispersive solid phase extraction cleanup (dSPE). QuEChERS methods, including an AOAC Official Method² and modifications to the methods, have been posted on the internet.³ These methods have several basic steps in common:

Step 1: Sample preparation and extraction – Commodities are uniformly homogenized. Acetonitrile solvent is added for a shake extraction. Salts, acids, and buffers may be added to enhance extraction efficiency and protect sensitive analytes. Surrogate standards can be added to monitor extraction efficiencies.

Step 2: Extract cleanup – A subsample of solvent extract is cleaned up using dSPE, a key improvement incorporated in the QuEChERS technique. Small polypropylene centrifuge tubes are prefilled with precise weights of MgSO₄ and SPE adsorbents to remove excess water and unwanted contaminants from the extracted samples. After agitation and centrifugation, the cleaned extracts are ready for analysis by a variety of techniques.⁴

Step 3: Sample analysis – Samples may be pH adjusted to protect sensitive pesticides and/or solvent-exchanged to improve analysis by either GC/MS or LC/MS. Internal standards can be added.

QuEChERS methods are convenient, rugged methods that simplify extract cleanup, reduce material costs, and improve sample throughput. Here we demonstrate the effectiveness of QuEChERS sample cleanup using a multiresidue analysis of pesticides on strawberries.

Experimental

Strawberry extracts were prepared, spiked, and dSPE treated according to Table I. One microliter splitless injections of the extracts were performed by a Shimadzu AOC-20i autosampler using “mid” injection speed into a Shimadzu QP-2010 Plus GC-MS system operated under the conditions in Table II.

Results and Discussion

Primary and secondary amine exchange material (PSA) is the base sorbent used for dSPE cleanup of QuEChERS fruit and vegetable extracts because it removes many organic acids and sugars that might act as instrumental interferences.

A pesticide-spiked strawberry extract (200ng/mL) subjected to dSPE with PSA was used to generate one-point calibration curves. Spiked strawberry extracts subjected to additional dSPE sorbents were analyzed and the results versus PSA dSPE are shown as percent recoveries in Table III. C18 is suggested for use when samples might contain fats; not an issue for a strawberry extract, but it was important to verify that gross losses of more hydrophobic pesticides (e.g., endrin and DDT) would not occur. Graphitized carbon black (GCB) is used to remove pigments, and when treated, the pink/red strawberry extract became clear. However, GCB can also have a negative effect on certain pesticides, especially those that can assume a planar shape like chlorothalonil and thiabendazole.



Restek dSPE products, available in a variety of standard sizes, make QuEChERS even simpler. The centrifuge tube format contains magnesium sulfate (to partition water from organic solvent) and a choice of SPE sorbents, including PSA (to remove sugars and fatty acids), C18 (to remove nonpolar interferences such as fats), and GCB (to remove pigments and sterols). Custom products also are available by request. If you are frustrated by the time and cost involved with your current approach to pesticide sample cleanup, we suggest you try this simple and economical new method.

References

1. Michelangelo Anastassiades, Steven J. Lehotay, Darinka Štajnbaher, Frank J. Schenck. "Fast and Easy Multiresidue Method Employing Acetonitrile Extraction/Partitioning and Dispersive Solid-Phase Extraction for the Determination of Pesticide Residues in Produce." *J. AOAC International*, 2003, vol. 86(22), pp.412-431.
2. AOAC Official Method 2007.01, "Pesticide Residues in Foods by Acetonitrile Extraction and Partitioning with Magnesium Sulfate."
3. <http://www.quechers.com/>
4. Schenck, F.J., *SPE Cleanup and Analysis of PPB Levels of Pesticides in Fruits and Vegetables*. Florida Pesticide Residue Workshop, 2002. References not available from Restek

Table I QuEChERS extraction and cleanup procedure for pesticides from strawberries.

Sample preparation and extraction

Sample:	10g of strawberries were homogenized and placed in a 50mL PTFE centrifuge tube (cat.# 26227)
Solvent:	10mL of acetonitrile were added to homogenate Shake for 1 minute, until uniform
Salts:	4.0g MgSO ₄ (powder or granular) 1.0g NaCl 1.0g trisodium citrate dihydrate 0.5g disodium hydrogencitrate sesquihydrate Salts were added and vigorously shaken for 1 minute. Sample was centrifuged and the supernatant removed for cleanup. Pesticides standards (200ng/mL) were spiked in at this point.

(cat.# 26213)

Sample extract cleanup

QuEChERS tubes:	1mL of supernatant from the previous step was placed into several 2mL polypropylene centrifuge tubes, each containing one of the following adsorbent mixes: A. 50mg PSA + 150mg MgSO ₄ (cat.# 26124) B. 50mg PSA + 150mg MgSO ₄ + 50mg C18 (cat.# 26125) C. 50mg PSA + 150mg MgSO ₄ + 50mg GCB (cat.# 26123)
Cleanup:	Samples were shaken with the adsorbents for 30 seconds (carbon for 2 minutes), then centrifuged to produce a clear supernatant for GC/MS analysis.
Internal standard:	Pentachloronitrobenzene in a formic acid solution, pH 5.

PSA—primary and secondary amine exchange material.
GCB—graphitized carbon black

Table II Instrument conditions.

Column:	Rtx [®] -CLPesticides2, 20m, 0.18mm ID, 0.14μm (cat.# 42302)
Sample:	custom pesticide mix, 200μg/mL each pesticide, internal standards: 8140-8141 ISTD, 1000μg/mL (cat.# 32279), 508.1 ISTD 100μg/mL (cat.# 32091), triphenylphosphate 1000μg/mL (cat.# 32281)
Inj.:	1.0μL splitless (hold 1 min.)
Inj. temp.:	250°C
Carrier gas:	helium
Flow rate:	constant linear velocity @ 40cm/sec
Oven temp.:	40°C (hold 1 min.) to 320°C @ 12°C/min.
Det:	Shimadzu GCMS-QP2010 Plus
Transfer line temp.:	300°C
Ionization:	Electron ionization
Mode:	Selected ion monitoring



did you know?

Multiple sorbents are used to extract different types of interferences.

MgSO ₄	removes excess water
PSA*	removes sugars, fatty acids, organic acids, and anthocyanine pigments
C18	removes nonpolar interferences
GCB**	removes pigments, sterols, and nonpolar interferences

*PSA—primary and secondary amine exchange material.

**GCB—graphitized carbon black

Table III Pesticide percent recoveries in strawberry extracts treated with C18 or GCB dSPE, relative to PSA only.

Rt (min.)	pesticide	CAS Number	action/use	classification	C18*	GCB**
9.50	dichlorvos	62-73-7	insecticide	organophosphorus	111	116
9.67	methamidophos	10265-92-6	insecticide	organophosphorus	105	107
11.75	mevinphos	7786-34-7	insecticide	organophosphorus	112	130
12.02	<i>o</i> -phenylphenol	90-43-7	fungicide	unclassified	106	97
12.14	acephate	30560-19-1	insecticide	organophosphorus	128	147
13.89	omethoate	1113-02-6	insecticide	organophosphorus	120	119
14.74	diazinon	333-41-5	insecticide	organophosphorus	108	127
14.98	dimethoate	60-51-5	insecticide	organophosphorus	124	151
15.69	chlorothalonil	1897-45-6	fungicide	organochlorine	125	13
15.86	vinclozolin	50471-44-8	fungicide	organochlorine	102	98
16.21	metalaxyl	57837-19-1	fungicide	organonitrogen	105	117
16.28	carbaryl	63-25-2	insecticide	carbamate	114	111
16.60	malathion	121-75-5	insecticide	organophosphorus	124	160
16.67	dichlofluanid	1085-98-9	fungicide	organohalogen	122	103
17.51	thiabendazole	148-79-8	fungicide	organonitrogen	88	14
17.70	captan	133-06-2	fungicide	organochlorine	88	91
17.76	folpet	133-07-3	fungicide	organochlorine	108	63
18.23	imazalil	35554-44-0	fungicide	organonitrogen	115	95
18.39	endrin	72-20-8	insecticide	organochlorine	104	101
18.62	myclobutanil	88671-89-0	fungicide	organonitrogen	119	114
19.07	4,4-DDT	50-29-3	insecticide	organochlorine	102	95
19.22	fenhexamid	126833-17-8	fungicide	organochlorine	118	77
19.40	propargite 1	2312-35-8	acaricide	organosulfur	110	95
19.43	propargite 2	2312-35-8	acaricide	organosulfur	121	114
19.75	bifenthrin	82657-04-3	insecticide	pyrethroid	106	81
20.04	dicofol	115-32-2	acaricide	organochlorine	98	54
20.05	iprodione	36734-19-7	fungicide	organonitrogen	118	90
20.21	fenpropathrin	39515-41-8	insecticide	pyrethroid	113	96
21.32	<i>cis</i> -permethrin	52645-53-1	insecticide	pyrethroid	106	65
21.47	<i>trans</i> -permethrin	51877-74-8	insecticide	pyrethroid	109	71
23.74	deltamethrin	52918-63-5	insecticide	pyrethroid	97	52

*50mg PSA, 50mg C18, **50mg PSA, 50mg GCB

$$\% \text{ recovery} = \frac{\text{RRF C18 or GCB}}{\text{RRF PSA}} \times 100$$

Rtx®-CLPesticides2 (proprietary Crossbond® phase)

- Application-specific columns for organochlorine pesticides and herbicides.
- Low bleed—ideal for GC/ECD or GC/MS analyses.
- Baseline separations in less than 10 minutes.

ID	df (µm)	temp. limits	length	cat. #
0.18mm	0.14	-60 to 310/330°C	20-meter	42302



Q-sep™ 3000 Centrifuge for QuEChERS

- Meets requirements of AOAC and European QuEChERS methodology.
- Supports 50mL, 15mL, and 2mL centrifuge tubes.
- Small footprint requires less bench space.
- Safe and reliable—UL, CSA, and CE approved, 1-year warranty.

Description	qty.	cat.#
Q-sep 3000 Centrifuge, 110V	ea.	26230
Q-sep 3000 Centrifuge, 220V	ea.	26231
Replacement Accessories		
50mL Tube Carrier for Q-sep 3000 Centrifuge	2-pk.	26232
50mL Conical Tube Insert for Q-sep 3000 Centrifuge	6-pk.	26235
4-Place Tube Carrier for Q-sep 3000 Centrifuge	2-pk.	26233
2mL Tube Adaptors for Q-sep 3000 Centrifuge	4-pk.	26234



QuEChERS SPE Tubes

Quick, Easy, Cheap, Effective, Rugged, and Safe, the QuEChERS (“catchers”) method is a fast, simple, and effective alternative to conventional sample prep for multiresidue pesticide analysis. Restek Q-sep™ products make QuEChERS even simpler. All extraction salts, adsorbents, and sample tubes are included—no specialized equipment or glassware is required.



cat. # 26123



cat. # 26124



cat. # 26227

cat. # 26214



cat. # 26125



cat. # 26126

Free Sample Packs Available!

To receive your free sample pack, add -248 to the item number. (One sample per customer.)

Description	Material	Methods	qty.	cat#
50mL Centrifuge Tubes for Sample Extraction				
Q110	4g MgSO ₄ , 1g NaCl, 1g trisodium citrate dihydrate, .5g disodium hydrogencitrate sesquihydrate	European EN 15662	50-pk.	26213
Q150	6g MgSO ₄ , 1.5g NaOAc	AOAC 2007.1	50-pk.	26214
Empty 50mL Centrifuge Tube	—	European EN 15662, AOAC 2007.1	25-pk.	26227
2mL Micro-Centrifuge Tubes for dSPE (clean-up of 1mL extract)				
Q210	150mg MgSO ₄ , 25mg PSA	European EN 15662	100-pk.	26215
Q211	150mg MgSO ₄ , 25mg PSA, 25mg C18	—	100-pk.	26216
Q212	150mg MgSO ₄ , 25mg PSA, 2.5mg GCB	European EN 15662	100-pk.	26217
Q213	150mg MgSO ₄ , 25mg PSA, 7.5mg GCB	European EN 15662	100-pk.	26218
Q250	150mg MgSO ₄ , 50mg PSA	AOAC 2007.1	100-pk.	26124
Q251	150mg MgSO ₄ , 50mg PSA, 50mg C18	AOAC 2007.1	100-pk.	26125
Q253	150mg MgSO ₄ , 50mg PSA, 50mg GCB	—	100-pk.	26123
Q252	150mg MgSO ₄ , 50mg PSA, 50mg C18, 50mg GCB	AOAC 2007.1	100-pk.	26219
15mL Centrifuge Tubes for dSPE (clean-up of 6mL extract)				
Q350	1200mg MgSO ₄ , 400mg PSA	AOAC 2007.1	50-pk.	26220
Q351	1200mg MgSO ₄ , 400mg PSA, 400mg C18	AOAC 2007.1	50-pk.	26221
Q352	1200mg MgSO ₄ , 400mg PSA, 400mg C18, 400mg GCB	AOAC 2007.1	50-pk.	26222
Q370	900mg MgSO ₄ , 150mg PSA	European EN 15662	50-pk.	26223
Q371	900mg MgSO ₄ , 150mg PSA, 15mg GCB	European EN 15662	50-pk.	26224
Q372	900mg MgSO ₄ , 150mg PSA, 45mg GCB	European EN 15662	50-pk.	26225
Q373	900mg MgSO ₄ , 150mg PSA, 150mg C18	—	50-pk.	26226
Q374	900mg MgSO ₄ , 300mg PSA, 150mg GCB	—	50-pk.	26126

Prepare samples more quickly, easily, and cost-effectively with QuEChERS.

	Mini-Luke or Modified Luke Method	QuEChERS	Savings with QuEChERS
Estimated time to process 6 samples (min.)	120	30	4x faster
Solvent used (mL)	60-90	10	6-9x less solvent
Chlorinated waste (mL)	20-30	0	Safer, cheaper, greener
Glassware/specialized equipment	capacity for 200mL, quartz wool, funnel, water bath or evaporator	none	Ready-to-use



Visit www.restek.com/quechers for detailed technical literature and a complete line of QuEChERS products.

PATENTS & TRADEMARKS
Restek patents and trademarks are the property of Restek Corporation. Other trademarks appearing in Restek literature or on its website are the property of their respective owners.



Lit. Cat.# GNAN1097A

© 2009 Restek Corporation.

Restek U.S. • 110 Benner Circle • Bellefonte, PA 16823 • 814-353-1300 • 800-356-1688 • fax: 814-353-1309 • www.restek.com

Restek France • phone: +33 (0)1 60 78 32 10 • fax: +33 (0)1 60 78 70 90 • e-mail: restek@restekfrance.fr

Restek GmbH • phone: +49 (0)6172 2797 0 • fax: +49 (0)6172 2797 77 • e-mail: info@restekgmbh.de

Restek Ireland • phone: +44 (0)2890 814576 • fax: +44 (0)2890 814576 • e-mail: restek@restek.com

Restek Japan • phone: +81 (3)6459 0025 • fax: +81 (3)6459 0025 • e-mail: ryosei.kanaguchi@restek.com

Thames Restek U.K. LTD • phone: +44 (0)1494 563377 • fax: +44 (0)1494 564990 • e-mail: sales@thamesrestek.co.uk

